

February 7, 2008
BA-TH/93-147
DPUR 65

A Coherent Understanding of Solvable Models for Quantum Measurement Processes

Hiromichi NAKAZATO⁽¹⁾ & Saverio PASCAZIO⁽²⁾

⁽¹⁾Department of Physics, University of the Ryukyus, Okinawa 903-01,
Japan

⁽²⁾Dipartimento di Fisica, Università di Bari
and Istituto Nazionale di Fisica Nucleare, Sezione di Bari
I-70126 Bari, Italy

Abstract

By making use of Schwinger's oscillator model of angular momentum, we put forward an interesting connection among three solvable Hamiltonians, widely used for discussions on the quantum measurement problem. This connection implies that a particular macroscopic limit has to be taken for these models to be physically sensible.

PACS: 03.65.Fd; 03.80.+r; 42.52.+x; 03.65.Bz

The quantum measurement problem [1] [2] has been a central issue in the foundations and interpretation of quantum mechanics, and is profoundly related to the investigation of the internal consistency of quantum theory. In order to find a resolution within the quantum mechanical framework, one usually describes a quantum measurement process in the following way: An elementary particle Q , on which the measurement is performed, interacts with a detection system D , which is considered to be made up of N elementary constituents. Generally, it is not easy to handle such a complicated system like $Q + D$, however, it is also true that several solvable models of measurement processes are now at hand. Among these are the Cini [3] and Coleman-Hepp [4] Hamiltonians and its modified version [5], which were all proposed for consideration of quantum measurement processes, and the Jaynes-Cummings [6] model, that describes the interaction between a two-level atom and a single electromagnetic mode in a cavity.

Although these three solvable models were proposed independently, a great deal of similarity exists among them. Indeed, we can establish several links among them in certain limits [7] [5]. This is both interesting and rather curious. The purpose of this letter is to gain a more comprehensive understanding on such models for quantum measurement processes, and in particular on their macroscopic limits, by making use of Schwinger's double-oscillator method [8] [9]. We stress that in all the above-mentioned cases, both the macroscopic system and the measurement process are described according to the quantum mechanical laws, and no "classical" behavior is postulated.

Let us first briefly review Schwinger's method. Schwinger's construction of the algebra of angular momentum originates from two independent harmonic oscillators, henceforth conventionally denoted as type $+$ and $-$. The creation and annihilation operators $a_{\pm}^{\dagger}, a_{\pm}$ obey the standard boson commutation relations:

$$\begin{aligned} [a_{\pm}, a_{\pm}^{\dagger}] &= 1, \\ [\mathcal{N}_{\pm}, a_{\pm}] &= -a_{\pm}, \\ [\mathcal{N}_{\pm}, a_{\pm}^{\dagger}] &= a_{\pm}^{\dagger}, \end{aligned} \tag{1}$$

where $\mathcal{N}_{\pm}$ are the number operators. All commutators of operators of different oscillators are assumed to vanish: This is what is meant by *independent* oscillators.

One can construct the eigenstates of $\mathcal{N}_{\pm}$ by acting with $a_{\pm}^{\dagger}$ on the *vacuum* state, defined by

$$a_{\pm}|0, 0\rangle = 0. \tag{2}$$

For instance, one has

$$|n_{+}, n_{-}\rangle = \frac{1}{\sqrt{n_{+}!}} \frac{1}{\sqrt{n_{-}!}} (a_{+}^{\dagger})^{n_{+}} (a_{-}^{\dagger})^{n_{-}} |0, 0\rangle. \tag{3}$$

Schwinger's interesting and brilliant idea was to define [we set $\hbar = 1$ in eqs.(4)-(9)]

$$J_{\pm} \equiv a_{\pm}^{\dagger} a_{\mp}, \tag{4}$$

$$J_3 \equiv \frac{1}{2} (a_{+}^{\dagger} a_{+} - a_{-}^{\dagger} a_{-}). \tag{5}$$

The above operators obey the standard commutation relations for angular momentum:

$$\begin{aligned} [J_3, J_{\pm}] &= \pm J_{\pm}, \\ [J_+, J_-] &= 2J_3. \end{aligned} \quad (6)$$

Moreover

$$\mathbf{J}^2 = J_3^2 + \frac{1}{2}(J_+J_- + J_-J_+) = \frac{\mathcal{N}_{\text{tot}}}{2} \left(\frac{\mathcal{N}_{\text{tot}}}{2} + 1 \right), \quad (7)$$

where $\mathcal{N}_{\text{tot}} \equiv \mathcal{N}_+ + \mathcal{N}_-$. By identifying

$$\begin{aligned} n_+ &\leftrightarrow j + m, & \frac{n_+ + n_-}{2} &\leftrightarrow j, \\ n_- &\leftrightarrow j - m, & \frac{n_+ - n_-}{2} &\leftrightarrow m, \end{aligned} \quad (8)$$

where j, m are the eigenvalues of $\mathbf{J}^2$ and J_3 , respectively, one can construct the familiar simultaneous eigenstates of $\mathbf{J}^2$ and J_3 :

$$|j, m\rangle \equiv \frac{1}{\sqrt{(j+m)!}} \frac{1}{\sqrt{(j-m)!}} (a_+^\dagger)^{j+m} (a_-^\dagger)^{j-m} |0\rangle. \quad (9)$$

Let us now apply Schwinger's procedures to show how it is possible to obtain the Coleman-Hepp Hamiltonian [4] from Cini's [3]. This will have interesting spin-offs: The conditions under which the two Hamiltonians can be identified [7] will suggest a link with the famous Jaynes-Cummings Hamiltonian [6], widely used in laser theory and recently utilized to discuss fundamental issues in quantum mechanics [5] [10]. Here and in the following we shall restrict our attention to the interaction term of the total Hamiltonian, and shall comment on the role of the free part whenever necessary.

In the model proposed by Cini, a quantum particle interacts with a detector D , made up of N particles, each of which is assumed to have only two possible states, say a ground state ω_- and an excited or ionized state ω_+ . The interaction Hamiltonian is taken to be

$$H_C = g (a_-^\dagger a_+ + a_+^\dagger a_-), \quad (10)$$

where g is a coupling constant and $a_\pm^\dagger, a_\pm$ are the creation and annihilation operators for the states $\omega_\pm$. We are here neglecting a (trivial) $\frac{1}{2}(1 + \tau_3)$ factor, whose only effect is to select which of the two states of the Q particle interacts with the detector.

By making use of eq.(4) we can write

$$H_C = g (J_+ + J_-) = g J_1. \quad (11)$$

Notice that H_C is essentially expressed as one of the group elements of angular momentum.

The Coleman-Hepp (CH) or AgBr Hamiltonian [4] [11] describes the interaction between a particle Q and a one-dimensional N -spin array (D -system). One can think, for instance, of a linear emulsion of AgBr molecules, the *down* state corresponding to

the undivided molecule, and the *up* state corresponding to the dissociated molecule (Ag and Br atoms). The particle and each molecule interact via a spin-flipping local potential, according to the following interaction Hamiltonian

$$H_{\text{CH}} = \sum_{n=1}^N V(\hat{x} - x_n) \sigma_1^{(n)}, \quad (12)$$

where $\hat{x}$ is the position of the particle, V is a real potential, x_n ($n = 1, \dots, N$) are the positions of the scatterers in the array and $\sigma_1^{(n)}$ is the Pauli matrix acting on the n th site. This Hamiltonian is a nice model of a typical measurement process and can be solved exactly if the Q particle is “ultrarelativistic”, namely if its free Hamiltonian is written as $H_Q = c\hat{p}$, $\hat{p}$ being the particle’s momentum. In such a case, the S -matrix is readily computed as

$$S^{[N]} = \exp \left(-i \frac{V_0 \delta}{\hbar c} \sum_{n=1}^N \sigma_1^{(n)} \right) = \exp \left(-i \frac{V_0 \delta}{\hbar c} N \Sigma_1^{(N)} \right), \quad (13)$$

where $V_0 \delta \equiv \int_{-\infty}^{\infty} V(x) dx$ is the integrated strength of the potential, $q \equiv \sin^2(V_0 \delta / \hbar c)$ is viewed as the probability of dissociating one AgBr molecule, and we defined the average spin

$$\Sigma_j^{(N)} = \frac{1}{N} \sum_{n=1}^N \sigma_j^{(n)}, \quad j = 1, 2, 3. \quad (14)$$

If the initial D -state is taken to be the ground state $|0\rangle_N$ (N spins down), and the initial Q -state is a plane wave, the evolution is

$$S^{[N]} |p, 0\rangle_N = \sum_{j=0}^N \binom{N}{j}^{1/2} (-i\sqrt{q})^j \left(\sqrt{1-q} \right)^{N-j} |p, j\rangle_N, \quad (15)$$

where $|j\rangle_N$ is the (symmetric) state with j dissociated molecules, and we have used the notation $|p, j\rangle_N = |p\rangle |j\rangle_N$. The right hand side in eq.(15) is a generalized [SU(2)] coherent state. Notice the (square root of the) binomial distribution in the above equation.

Observe that the operators $N \Sigma_j^{(N)}$ form a unitary representation of SU(2), and therefore satisfy the commutation relations (6). Notice that in this representation, the Cini Hamiltonian (11) is expressed as

$$H_C = g \left(N \Sigma_+^{(N)} + N \Sigma_-^{(N)} \right) = g N \Sigma_1^{(N)} = g \sum_{n=1}^N \sigma_1^{(n)}. \quad (16)$$

This expression is very similar to the CH Hamiltonian (12), and indeed, H_C appears as a particular case of H_{CH} : Consider either the situation in which the spins are all placed at the same position

$$x_n \equiv x_0 = 0, \quad \forall n = 1, \dots, N \quad (17)$$

or the (similar) situation of an “average” potential over the positions of the scatterers x_n (replace $V(\hat{x} - x_n)$ with its average, say $\bar{V}(\hat{x})$, and call the latter $V(\hat{x})$ again. Both cases have been analyzed in Ref. [7]: H_C and H_{CH} coincide if one takes $V(\hat{x}) = \text{constant} = g$. It is necessary to observe, in this context, that unlike CH, Cini’s Hamiltonian does not contain a free part, involving the Q -particle coordinates, so that one is forced to assume that the Q and D systems are in contact for a certain “interaction time” τ . In this sense, the CH Hamiltonian appears more self-contained, because the interaction time is naturally given by $\tau = L/c$, where L is the total length of the spin array and c the Q -particle speed. Interestingly, we shall also see that τ cannot be arbitrarily chosen for H_C : This will shed light on Cini’s choice for τ .

In order to establish a link with the Jaynes-Cummings model, we need to consider the weak-coupling, $N \rightarrow \infty$ limit of the CH Hamiltonian [5]. Notice that H_{CH} is invariant under exchange of spins (molecules) in the array. Therefore, if we call $\mathcal{P}_N$ the group of permutations on $\{1, \dots, N\}$, we can restrict our attention to the $\mathcal{P}_N$ -invariant sector $\mathcal{H}_N$ of the bigger Hilbert space $\mathcal{H}_{\{N\}}$ of the N spins. In the following, we shall concentrate our analysis on the *symmetrized* case.

The weak-coupling, $N \rightarrow \infty$ limit of H_{CH} can be consistently taken only if we consider a modified version of the CH model, that enables us to take into account the possibility of energy exchange between the particle and the spin system. One writes the interaction part as

$$\begin{aligned} H'_{CH} &= \sum_{n=1}^N V(\hat{x} - x_n) \sigma_1^{(n)} \exp\left(i \frac{\omega}{c} \sigma_3^{(n)} \hat{x}\right) \\ &= \sum_{n=1}^N V(\hat{x} - x_n) \left[\sigma_+^{(n)} \exp\left(-i \frac{\omega}{c} \hat{x}\right) + \sigma_-^{(n)} \exp\left(+i \frac{\omega}{c} \hat{x}\right) \right], \end{aligned} \quad (18)$$

and adds the free Hamiltonian of the spin array $H_D = \frac{1}{2} \hbar \omega \sum_{n=1}^N (1 + \sigma_3^{(n)})$. The reasons for this modification and additional details can be found in Ref. [5]: It is remarkable that the model remains solvable provided that a “resonance condition” (implicitly assumed in writing the above expressions) is met, because in this way the energy acquired or lost by the Q -particle in every single interaction matches exactly the energy gap between the two spin states. In fact, there is a deep reason for this: It is practically equivalent to the rotating-wave approximation in optics.

The weak-coupling, $N \rightarrow \infty$ limit of H_{CH} can be consistently taken by performing a contraction on $SU(2)$ [12]: We start from a linear transformation of the generators of $SU(2)$

$$\begin{pmatrix} h_+ \\ h_- \\ h_3 \\ 1 \end{pmatrix} = \begin{pmatrix} N^{-1/2} & & & \\ & N^{-1/2} & & \\ & & 1 & N/2 \\ & & & 1 \end{pmatrix} \begin{pmatrix} N \Sigma_+^{(N)} \\ N \Sigma_-^{(N)} \\ N \Sigma_3^{(N)}/2 \\ \mathbf{1}^{(N)} \end{pmatrix}, \quad (19)$$

so that the commutation properties for $\mathbf{h}, 1$ are

$$[h_3, h_{\pm}] = \pm h_{\pm},$$

$$\begin{aligned}
[h_-, h_+] &= 1 - \frac{2}{N} h_3, \\
[\mathbf{h}, 1] &= 0,
\end{aligned} \tag{20}$$

and yield, in the $N \rightarrow \infty$ limit, the standard boson commutation relations:

$$\begin{aligned}
h_+ &= \frac{1}{\sqrt{N}} \sum_{n=1}^N \sigma_+^{(n)} = \sqrt{N} \Sigma_+^{(N)} \xrightarrow{N \rightarrow \infty} a^\dagger, \\
h_- &= \frac{1}{\sqrt{N}} \sum_{n=1}^N \sigma_-^{(n)} = \sqrt{N} \Sigma_-^{(N)} \xrightarrow{N \rightarrow \infty} a, \\
h_3 &= \frac{1}{2} \sum_{n=1}^N (1 + \sigma_3^{(n)}) = \frac{N}{2} (\mathbf{1}^{(N)} + \Sigma_3^{(N)}) \xrightarrow{N \rightarrow \infty} \mathcal{N} \equiv a^\dagger a.
\end{aligned} \tag{21}$$

Now assume again, for simplicity, that all spins are placed at the same position, so that the Hamiltonian (18) becomes

$$\begin{aligned}
H'_{\text{CH}} &= V(\hat{x}) \sum_{n=1}^N \left[\sigma_+^{(n)} \exp\left(-i\frac{\omega}{c}\hat{x}\right) + \sigma_-^{(n)} \exp\left(+i\frac{\omega}{c}\hat{x}\right) \right] \\
&= V(\hat{x}) \sqrt{N} \left[\exp\left(-i\frac{\omega}{c}\hat{x}\right) \frac{1}{\sqrt{N}} \sum_{n=1}^N \sigma_+^{(n)} + \exp\left(+i\frac{\omega}{c}\hat{x}\right) \frac{1}{\sqrt{N}} \sum_{n=1}^N \sigma_-^{(n)} \right] \\
&\xrightarrow{N \rightarrow \infty, V\sqrt{N}=u} u(\hat{x}) \left[a^\dagger \exp\left(-i\frac{\omega}{c}\hat{x}\right) + a \exp\left(+i\frac{\omega}{c}\hat{x}\right) \right] = H_{\text{JC}}.
\end{aligned} \tag{22}$$

The connection with a “maser” system is manifest: In the weak-coupling, $N \rightarrow \infty$ limit we obtain the Jaynes-Cummings (JC) [6] Hamiltonian. (Strictly speaking, the JC Hamiltonian differs from the above one because it contains terms of the type $\tau_\pm$, instead of $\exp(\pm i\frac{\omega}{c}\hat{x})$, $\tau_\pm$ being the raising/lowering operators for a two-level system. In the case we are considering the Q particle has a continuous spectrum, and can exchange an arbitrary number of quanta of energy $\hbar\omega$. Clearly, this difference is not important for our analysis.)

It is necessary to observe that the above limit is taken by keeping $V(\hat{x})\sqrt{N} = u(\hat{x})$ finite, while sending N to infinity. This is just a *consequence* of the (physically appealing) requirement that the average number of dissociated molecules, namely the quantity qN , with $\sqrt{q} \simeq V_0\delta/\hbar c = \int V(y)dy/\hbar c$ [see the definition after eq.(13)], is kept finite in the $N \rightarrow \infty$ limit.

Incidentally, notice that in this limit the free Hamiltonian of the spin array, introduced after eq.(18), yields just the free Hamiltonian of a single-mode electromagnetic cavity:

$$H_D = \frac{1}{2} \hbar\omega \sum_{n=1}^N (1 + \sigma_3^{(n)}) \xrightarrow{N \rightarrow \infty} \hbar\omega \mathcal{N} = \hbar\omega a^\dagger a = H_D^{\text{JC}}. \tag{23}$$

The N -spin system behaves, in the above-mentioned limit, as a “cavity”, in which boson-like excitations (collective modes) can be created, as a consequence of the interaction with the Q -particle.

The S -matrix of the JC Hamiltonian is

$$S = \exp \left[-i \frac{u_0 \delta}{\hbar c} \left(a^\dagger \exp \left[-i \frac{\omega}{c} \hat{x} \right] + a \exp \left[i \frac{\omega}{c} \hat{x} \right] \right) \right], \quad (24)$$

where $\int u(x) dx = u_0 \delta$, and we assumed, for simplicity and without loss of generality, that δ is the same quantity used in eq.(13). If we take the initial state to be $|p, 0\rangle = |p\rangle |0\rangle$, where $|0\rangle$ is the ground state of the maser cavity, the evolution is

$$S|p, 0\rangle = e^{-\bar{\kappa}/2} \sum_{j=0}^{\infty} \frac{(-i\sqrt{\bar{\kappa}})^j}{\sqrt{j!}} |p-j, \frac{\hbar\omega}{c}, j\rangle, \quad \bar{\kappa} = \left(\frac{u_0 \delta}{\hbar c} \right)^2, \quad (25)$$

where $|j\rangle$ is the number state of the cavity and $\bar{\kappa}$ is the average number of boson excitations in the cavity. This reduces to the $N \rightarrow \infty$, $qN \equiv \bar{\kappa}$ finite limit of eq.(15) when we neglect the energy differences between spin states, i.e., $\omega = 0$. [The general case, with $\omega \neq 0$, has been analyzed in Ref. [5] and yields a final state that is essentially and conceptually similar to (15).]

We wish to stress that the connection between H'_{CH} and H_{JC} has been proven under the condition that all spin were at the same position in space, like in eq.(17). The fact that such a condition is unnecessary was already stressed in Ref. [5], and in fact a rigorous derivation, that does not make use of (17) has recently been found [13].

We emphasized before that the $N \rightarrow \infty$ limit was taken by keeping $\sqrt{qN} = (V_0 \delta / \hbar c) \sqrt{N} = (\int V(y) dy / \hbar c) \sqrt{N}$ finite. Since the $Q + D$ system is isolated, its total energy is conserved, so that $qN \hbar \omega$, the energy necessary to provoke the dissociation of qN AgBr molecules (namely, the average energy “stored in the detector”), is also the energy lost by the Q particle. Such an energy *must* be finite, because if it were not finite we would encounter a pathological situation of the model, in which the Q particle has a negative energy: The above limit appears therefore as a *forced choice from the physical point of view*. Notice also that, in the above limit, the binomial statistics in eq.(15) yields the Poisson statistics of eq.(25). Obviously, this could be accomplished *only* by keeping $qN = \bar{\kappa}$ finite. Returning to our AgBr model, the probability q of one spin flip or one AgBr molecule dissociation becomes very low, of order $O(N^{-1})$: In the macroscopic limit, the event of one dissociation becomes therefore a *rare event* in the statistical sense, for the AgBr model to make sense from a physical point of view.

In this context, observe that Cini redefined his coupling constant as

$$g \equiv \frac{g_0}{\sqrt{N}}, \quad (26)$$

so that $\tau_0 = \hbar / g \sqrt{N} = \hbar / g_0$, the time required to ionize the first particle of the initially discharged detector, becomes independent of N , the number of elementary constituents of his “detector”. Since g is nothing but our $V_0 \delta \propto \sqrt{q}$, Cini’s intuition appears as a *straight consequence of the macroscopicity of the detection system*. Indeed, according to the above discussion, one must require that $g^2 \propto q = O(N^{-1})$, for the model to be physically sensible.

There are also profound links between the limiting procedure considered in this letter and van Hove's " $\lambda^2 T$ " limit [14]. This is easily evinced from the following example: Set $x_n \equiv na$ in eq.(12) or (18) (constant spacing between adjacent potentials). The free part of the JC Hamiltonian is $H_Q = c\hat{p}$, so that the particle travels with constant speed c , and interacts with the detector for a time $T = Na/c$, where Na is the total length of the detector. Since the coupling constant $\lambda \equiv g \propto V_0\delta$, one gets $\lambda^2 T = g^2 Na/c \propto qN$. (Observe that the "lattice spacing" a , the inverse of which corresponds to a density in our 1-dimensional model, is kept constant in the above limit.) Note also that the hypothesis of photon-like dispersion-free Q particles ($H_Q = c\hat{p}$), typical of CH-like models, does not appear very restrictive: In similar cases, such as the JC model [10], in order to solve the equations of motion, it is often assumed that the particle has constant speed while travelling through the maser. Incidentally, we stress that this assumption is in line with our qN -finite ansatz: The energy lost by the particle during the interaction is negligible compared to its kinetic energy ($qN\hbar\omega \ll cp$).

We have mentioned some aspects of the quantum theory of measurement and stressed the importance of performing a full quantum mechanical analysis of the (macroscopic) "detector". The purpose of this work was to clarify the similarity among some model detectors, and illustrate that no "classical" behavior is to be postulated.

Many important topics, such as the problem of the loss of quantum coherence [15] and the general argument of "objectification" [16], namely the emergence of definite outcomes in a measurement process, have not been addressed. The problem of a consistent description of the interaction between a macroscopic and a microscopic object [17] is delicate and, in our opinion, still open. The unitarity of the quantum evolutions appears to many physicists as an insuperable problem, and has led some authors to conjecture that nonlinear terms should be added to the Schrödinger equation [18]. This is a logically consistent possibility that requires further investigation.

At the present stage, we feel that no solution, among those proposed, is fully satisfactory. Nevertheless, we are confident that further progress on these fundamental quantum mechanical problems will also come from "technical" analyses. Solvable models often provide excellent clues in this direction.

The authors acknowledge interesting remarks by Professors Y.G. Lu and L. Accardi. They thank the University of the Ryukyus Foundation and the Italian National Institute for Nuclear Physics (INFN) for the financial support. This work was also partially supported by the Grant-in-Aid for Scientific Research of the Ministry of Education, Science and Culture, Japan (03854017) and by Italian Research National Council (CNR) under the bilateral projects Italy-Japan n.91.00184.CT02 and n.92.00956.CT02.

References

- [1] J. von Neumann, *Die Mathematische Grundlagen der Quantenmechanik* (Springer Verlag, Berlin, 1932).
- [2] *Quantum Theory and Measurement*, edited by J.A. Wheeler and W.H. Zurek (Princeton University Press, 1983)
- [3] M. Cini, *Nuovo Cimento* **73B** (1983) 27.
- [4] K. Hepp, *Helv. Phys. Acta* **45** (1972) 237.
- [5] H. Nakazato and S. Pascazio, *Phys. Rev. Lett.* **70** (1993) 1; *Phys. Rev. A*, in print.
- [6] E.T. Jaynes and F.W. Cummings, *Proc. IEEE* **51** (1963) 89.
- [7] H. Nakazato and S. Pascazio, *Phys. Lett.* **156A** (1991) 386; *Phys. Rev.* **A45** (1992) 4355.
- [8] J. Schwinger, in *Quantum Theory of Angular Momentum*, edited by L.C. Biedenharm and H. Van Dam (Academic Press, 1965) p.229.
- [9] J.J. Sakurai, *Modern quantum mechanics*, edited by San Fu Tuan (Benjamin Cummings, California, 1985) p.182.
- [10] N.B. Narozhny, J.J. Sanchez-Mondragon and J.H. Eberly, *Phys. Rev.* **A23** (1981) 236; J. Krause, M.O. Scully and H. Walther, *Phys. Rev.* **A36** (1987) 4547; M.O. Scully and H. Walther, *Phys. Rev.* **A39** (1989) 5229; J. Gea-Banacloche, *Phys. Rev. Lett.* **65** (1990) 3385; S.J.D. Phoenix and P.L. Knight, *Phys. Rev.* **A44** (1991) 6023.
- [11] J.S. Bell, *Helv. Phys. Acta* **48** (1975) 93; S. Machida and M. Namiki, *Proceedings of the International Symposium on Foundations of Quantum Mechanics*, edited by S. Kamefuchi et al. (Physical Society of Japan, Tokyo, 1984) p.136; M. Namiki, *Found. Phys.* **18** (1988) 29; S. Kudaka, S. Matsumoto and K. Kakazu, *Prog. Theor. Phys.* **82** (1989) 665; M. Namiki and S. Pascazio, *Found. Phys. Lett.* **4** (1991) 203; C.P. Sun, “*Quantum dynamical model for wave function reduction in classical and macroscopic limits*”, preprint ITP.SB-93-14; K. Hiyama and S. Takagi, “*Generalized Coleman-Hepp model and quantum coherence*”, preprint (1993).
- [12] R. Gilmore, *Lie Groups, Lie Algebras, and Some of Their Applications* (John Wiley & Sons, New York, 1974) p.460; F.T. Arecchi, E. Courtens, R. Gilmore and H. Thomas, *Phys. Rev.* **A6** (1972) 2211.
- [13] L. Accardi, Y.G. Lu and S. Pascazio, in preparation.
- [14] L. van Hove, *Physica* **21** (1955) 517. See also E.B. Davies, *Commun. Math. Phys.* **39** (1974) 91; G.L. Sewell, *Quantum theory of collective phenomena*, (Clarendon Press, Oxford, 1986).

- [15] S. Machida and M. Namiki, Prog. Theor. Phys. **63** (1980) 1457; 1833; M. Namiki and S. Pascazio, Phys. Rev. **A44** (1991) 39; Phys. Reports, in print.
- [16] P. Busch, P.J. Lahti and P. Mittelstaedt, *The quantum theory of measurement* (Springer-Verlag, Berlin, 1991).
- [17] A.O. Caldeira and A.J. Leggett, Phys. Rev. **A31** (1985) 1057; F. Haake and D.F. Walls, Phys. Rev. **A36** (1987) 730; F. Haake and M. Żukowski, Phys. Rev. **A47** (1993) 2506.
- [18] G.C. Ghirardi, A. Rimini and T. Weber, Phys. Rev. **D34** (1986) 470; L. Bonci, P. Grigolini, R. Roncaglia and D. Vitali Phys. Rev. **A47** (1993) 3538.